

UTILIZATION OF AGUNG SEMERU BANANA PEEL EXTRACT AS NATURAL HAND SANITIZER

by Dwì Nur Rikhma Sari

Submission date: 30-Jun-2020 05:24PM (UTC-0700)

Submission ID: 1352001661

File name: Artikel_2020_-_Dwinur_-_Inggris_2kolom.pdf (2.56M)

Word count: 4003

Character count: 20969

UTILIZATION OF AGUNG SEMERU BANANA PEEL EXTRACT AS NATURAL HAND SANITIZER

Dwi Nur Rikhma Sari¹, Septarini Dian Anitasari², Suci Ratnasari³

¹²³ Biology Education, FP. MIPA IKIP PGRI Jember

*email: rikhmasari_dnr@ikipjember.ac.id

ABSTRACT

Hand sanitizers generally contain Ethyl Alcohol 62%, softener, moisturizer and anti-bacterial compounds such as tryclosan, glycerol, tannin, saponins and other antimicrobial agents. The purpose of this study is to test the effectiveness of Agung Semeru banana peel extract as a natural hand sanitizer to inhibit fungal growth of *Candida albicans*. The antifungal activity test is carried out using the disk diffusion method. The use of this method is shown to measure the diameter of the area of resistance that occurs around paper discs that already contain antifungi in accordance with the concentration in each treatment. The results of this study indicate that the Agung Semeru banana peel extract of Lumajang variety is effective as a natural Hand sanitizer, this result is shown when statistical tests use Kruskal-wallis Asymp values. Sig = 0'000 < α (0.05), where there are differences in treatment, it is proven at a concentration of 7% on average DHP which is 1,135 greater than the concentration that was not given extract or control that is equal to 1,200, the higher the treatment or concentration more and more inhibitory zones or clear zones will form.

Keywords: Hand sanitizer, Banana peel of Agung Semeru Lumajang, *Candida albicans*

INTRODUCTION

Infectious diseases in the digestive tract are caused by several factors, one of which is by pathogenic microorganisms such as *Candida albicans* which can indirectly enter the human digestive tract through the mouth (Rosenthal, 2005). *Candida albicans* is a type of yeast pathogenic fungus most found in the digestive tract, skin surface and in the female reproductive tract which often results in excessive vaginal discharge and odor (Hawkins, 2011; Zubier *et al.*, 2010). In addition, *Candida albicans* is also able to form biofilms that can carry out the process of host cell invasion and has the ability to be resistant to antifungal compounds (Kusumaningtyas, 2015). One effort to prevent diseases caused by *Candida albicans* is by providing antifungal which can inhibit and disrupt the growth of *Candida albicans* (Febriani, 2014).

Various studies have been carried out to reduce the prevalence of disease caused by patho-

genic microorganisms, one of which is by maintaining hand hygiene through washing hands using soap which is proven to reduce the number of microorganisms that cause disease (Rachmayanti, 2013). Along with the busyness of the community, various soap replacement innovations have been found such as hand sanitizers which contain antiseptic compounds and are used to kill pathogenic microorganisms in the hands and inhibit the growth and metabolic activity of microbial disease (Radji *et al.*, 2012).

Use for too long, the antiseptic liquid contained in the hand sanitizer fluid makes the skin dry, because it contains alcohol, so it needs the addition of moisturizer and emolient substances that make the skin stay soft. In addition, alcohol content as a basic ingredient in hand sanitizers causes skin and eye irritation (Retnosari, 2007) due to the highly flammable nature of alcohol (Dyer *et al.*, 2000). Re-

cent research revealed that the use of alcohol-based hand sanitizers from chemicals turned out to have a considerable impact on health, in addition to flammable alcohol-based hand sanitizers can also increase the risk of viral infections that trigger inflammation of the digestive tract (Fadhilah, 2017). Therefore we need other alternatives that are friendly and safe for the environment, especially the skin with the basic ingredients of natural hand sanitizer, one of which is by using the Agung Semeru banana peel extract of Lumajang variety.

Banana variety of Agung Semeru Lumajang is one type of banana found in Lumajang Regency, East Java. The characteristics of Agung Semeru banana varieties can be seen from the color of the stems (bright red), the formation of unique fruit, the number of tillers 1-2 tillers / clumps, in addition to the size of large fruit (around 19 cm) and length (33-36 cm), the number of comb 1-2 comb / bunch with a weight of 10-20 kg / bunch (Prahardini, 2010). According to Sari (2017) the potential content of phytochemical compounds found in peel of Agung Semeru banana Lumajang are phenolic, terpenoids, saponins and alkaloids. This is also supported by research by Ehiowemwenguan *et al.* (2014) which states that banana peels contain glycosides, alkaloids, saponins, tannins, flavonoids, terpenoids and phenols which can inhibit the growth of pathogenic microbes.

Various researches on the potential of Agung Semeru banana variety Lumajang peel extract have been widely carried out, one of which is used as a herbal skin cream formulation from Agung Semeru banana peel (Sari *et al.*, 2017), but no research has been done that utilizes Agung Semeru banana peel extract Lumajang varieties are natural

ingredients for making hand sanitizers. Based on this background, the researchers wanted to conduct a study entitled Utilization of Semeru Agung Banana Skin Extract Lumajang variety as a Natural Basic Material for Hand Sanitizer which will be tested on the growth of pathogenic microbes, *Candida albicans*.

METHODS

This research is an experimental study conducted at the Biology Laboratories of FPMIPA IKIP PGRI Jember which uses a Completely Randomized Design (CRD) with various concentrations of treatment 0%, 1%, 5%, 7% (Aponno *et al.*, 2014). Data obtained by direct observation and counting techniques. Observation was carried out to observe or find out whether Lumajang banana peel was effective as a natural hand sanitizer and counting was done to determine the diameter of the inhibitory zone of *Candida albicans* growth.

Tools and Materials. The tools used in this study are: a set of maceration tools, rotary evaporator, waterbath, autoclave, Laminar Air Flow (LAF), analytical scales, ose needles, incubators, petri dishes, test tubes and tube racks, measuring cups, Erlenmeyers, stirrers, and tube racks, sonicator baths, fresh dryers, stirrers. The materials used are: peel of Agung Semeru banana variety Lumajang, media *Potatoes Dextrose Agar* (PDA), spiritus, cotton, Aluminum foil, sterile distilled water, markers, label paper, 70% alcohol, tissue, wood paper, rubber bands, physiological salts and fungi *Candida albicans*.

Sterilisasi Alat dan Bahan. All tools and materials to be autoclaved are washed and dried first, Erlenmeyer's mouth, test tube and measuring

cup are covered with cotton, while the petri dish, tube rack is wrapped in wood paper, then put into the autoclave at 121°C and a pressure 1 atm with time for 1 hour (Manan and Kharisma, 2012)

Make of Peel Banana Extract. Semeru banana peel is cleaned and then cut into small pieces, acrated until slightly dry, after drying then mashed by way of ground and then sieved with a 10 mesh sieve to obtain a dry simplicia of Semeru banana skin variety Lumajang, after that weigh as much 150 g. Then the simplicia was put into Erlenmeyer and dissolved with 1125 mL of sterile Aquades with a ratio of 1: 7: 5. Then the solution is stirred until homogeneous and put into an ultrasonic clening bath with a frequency of 42 KHz and left for ± 2 hours (Fuadi, 2012). The extraction results are then filtered using filter paper to get pure liquid extract. The liquid extract was then put into a freshdryer for ± 1x24 hours (Sari *et al.*, 2018; Zakiyah *et al.*, 2017) to obtain a dry extract of Semeru banana peel. The skin extract was further diluted back into 4 concentrations namely 0%, 1%, 5%, 7%.

Make a Gel Hand Sanitizer basis Na-CMC. The heated water is put in a beaker (homogenizer), then enter the Na-CMC, let it rise for about 15 minutes, while stirring until it thickens slightly. Add propylene glycol, glycerin and distilled water, stir until it forms a gel. Then enter the banana peel extract dissolved in beaker glass and stir until homogeneous, mixed using low speed homogenizer until thick and homogeneous (Hasyim *et al.*, 2012).

Formulation Gel Of Banana Agung Semeru peel Extract. For the manufacture of the Agung Semeru banana peel extract formulation follows the standard. According to Hamzah *et al* (2007), making the standard formula of Hand Sani-

tizer with banana peel extract has a percent increase (%), namely: Na-CMC 5%; Glycerin 10%; Propylene-glycol 5% and a solution of 100 mL. So that from the formulation obtained by Agung Semeru banana peel extract with various concentrations of 0%, 1%, 5%, and 7% concentration.

Make PDA Media. A total of 20 g of PDA (*Potatos Dextrose Agar*) was dissolved in Erlenmeyer with 500 ml distilled water. Heated on a hot plate while stirring until the solution becomes homogeneous. Homogeneous medium was sterilized in an autoclave at 121°C, 2 atm pressure for 15 minutes. (Sari *et al.*, 2017).

Rejuvenation Of Fungus. The medium PDA which has been sterilized in an autoclave, is then poured into several test tubes that have been sterilized and tilted and then left to harden. Fungi colonies were taken from pure cultures that were available, carried out aseptically with an ose needle and etched on the media so that it was tilted and then incubated in an incubator (Rostinawati, 2009).

Determiration of Hand sanitizer anti-fungal activity. The antimicrobial activity test was carried out using the diffusion method, using the PDA. 1 ml suspense test microbes were inoculated in 10 ml PDA, then poured into a petri dish. Then the paper disk / disk paper that has been immersed in a gel solution mixed with banana peel extract with a concentration of 0%, 1%, 5%, and 7%, respectively, is placed in a petri dish which contains PDA and microbial suspense media. Incubated 2x24 hours at 37°C, then observed and measured the area of resistance formed (Sari *et al.*, 2018).

Technical Data Analysis. The parameters observed in this study are the diameter of the growth inhibition zone fungi *Candida albicans* (cm), then

the data will be tested using the *Kruskall Wallis* 5% test where before homogeneity testing and normality +test data are tested using SPSS version 23.

RESULTS AND DISCUSSIONS

Results

The results of the diameter of the inhibitory growth zone of *Candida albicans* at various concentrations of Agung Semeru banana peel extract as a natural Hand Sanitizer material using the *Kruskall Wallis* 5% showed that there is a significant difference (table 2). But based on the mean diameter of the growth inhibition zone shows that the greater the concentration of the Agung Semeru banana variety Lumajang peel extract, the greater the diameter of the growth inhibition zone *Candida albicans* (table 1 and Figure 2).

Table 1. The mean diameter of *Candida albicans* growth inhibition zones at various concentrations of Agung Semeru banana peel extract of Lumajang variety.

Concentration of Banana peel extract of Agung Semeru Variety Lumajang	Mean diameter of <i>Candida albicans</i> growth inhibition zones (cm)
0% (Kontrol)	1,200 ± 0,0000 ^a
1%	1,200 ± 0,0000 ^a
5%	1,208 ± 0,0023 ^b
7%	1,315 ± 0,0035 ^c

Table 2. *Kruskall-wallis test's*

	bening
Chi-Square	18.363
Df	3
Asymp.Sig.	.000

Description: Asymp Value. Sig = 0'000 < α (0.05) means that there is a difference between treatments for the diameter of the growth inhibition zone

Statistical testing of the utilization of a natural hand sanitizer was performed using the *Kruskall Wallis* level of 5% (table 2) due to the Homogeneity and Normality Test results that the diameter of the

Candida albicans growth inhibition zone data showed no data normal and not homogeneous.

The antifungal activity test of Agung Semeru banana peel extract as a natural Hand sanitizer was carried out on *Candida albicans* fungi, with different concentrations. The results of antifungal activity test of Agung Semeru Hand banana peel extract natural sanitizer can be seen in Figure 2, where there are differences in inhibition zone diameter in each treatment.

Table 3. Formulation of Hand Sanitizer

Formulation Gel of Hand Sanitizer	Concentration Of Banana Peel extract (%)			
	0	1	5	7
Extract (g)	0	0,25	1,25	2,25
Na-CMC (g)	1,25	1,25	1,25	1,25
Glycerin (g)	2,5	2,5	2,5	2,5
Propilen-glikol (g)	1,25	1,25	1,25	1,25
Aquadest (mL)	25	25	25	25

The formulation of hand sanitizer made from Agung Semeru banana peel extract (table 3) produced will then be organoleptically tested with all concentration treatments (1%, 5% 7%) with the result in the form of pH 6 (adjusted to the pH on the market), has forms such as gel / semisloid, clear brown color due to Agung Semeru banana peel extract is slightly brownish in color and has a distinctive aroma of banana peel because it is not added with synthetic aroma. Based on observations for 4 weeks, there was no change in the hand sanitizer, so the gel base formulation that has been made has been stable enough to cause no chemical changes in the antiseptic gel component. However, storage at 37⁰C shows that the liquid gel has a younger oxidation as indicated by the change in color and the change in the shape of the gel to be more liquid, whereas at cold temperatures causes the gel formula to be more stable.




Figure 1. Test results of antifungal activity of Agung Semeru banana peel extract as a natural Hand sanitizer against *Candida albicans* fungi at each treatment concentration 0% no clear zone, 1% concentration no clear zone, 5% concentration in petri dish (C) = 1,21 cm (clear zone), and 7% concentration in petri dishes (B) averaged = 1.32 cm (clear zone), in petri dishes (C) averaged = 1.315 cm (clear zone).

The antifungal activity test is carried out using the disk diffusion method which is marked by the formation of a clear zone around the disc paper. The use of this method is shown to measure the diameter of the area of resistance that occurs around paper discs that already contain antifungi in accordance with the concentration in each treatment. Agung Semeru banana peel extract as a natural Hand sanitizer was tested on *Candida albicans* using PDA (Potatos Dextrose Agar) and 1.2 cm diameter paper discs on petri dishes, shown in Figures 1 and 2.

Based on analysis results and concentration graphs (Figure 2) shows that the higher the concentration, the greater the diameter of the inhibition zone that is impossible to inhibit the *Candida albicans*. A concentration of 1% (1.2 ± 0.0^a) showed no

significant difference with a concentration of 0% (1.2 ± 0.0^a) and a concentration of 7% (1.315 ± 0.0035^c) was the greatest concentration.

Discussions

Based on the results of statistical tests using Kruskal-wallis in the antifungal test of the banana peel extract of Agung Semeru variety Lumajang as a natural hand sanitizer has the effectiveness to inhibit the *Candida albicans*, this is indicated by the presence of inhibition zones or clear zones formed in the area around disc paper (Figure 1).




Figure 2. Diameter of growth inhibition zone (cm) Agung Semeru banana variety Lumajang peel extraction against *Candida albicans* at various concentration treatments

Based on the average statistical test, there was a significant difference between treatments even though the 1% concentration did not differ from the control concentration of 0% (figure 2, table 1). This is probably due to the soaking of paper disks that are too fast so that it affects the inhibitory growth of *Candida albicans*. The results of the average inhibition zone diameter is at a concentration of 7% can increase the inhibition of fungi, as evidenced at an average concentration of 7% DHP which is 1,135 cm greater than the concentration not given extract or control that is equal to 1,200 cm. So this shows that the higher concentration of banana peel extract

as a hand sanitizer shows the greater inhibitory diameter results (Figure 2). This shows that the higher the concentration of banana peel extract as a hand sanitizer, the greater the diameter of the zone of inhibitory growth of pathogenic microbes (Kholifah, 2017).

Even though the addition of the Agung Semeru banana peel did show a significant difference between treatments, the mean diameter of the *Candida albicans* growth inhibition zone produced at various concentrations (table 1) did not show too much difference. Agung Semeru banana peel extract contains secondary metabolite compounds such as are phenolic, terpenoids, saponins and alkaloids (Sari, 2017). The results of the average diameter of the *Candida albicans* growth inhibition zone produced are not so visible that the difference is quite far, this is likely influenced by several factors such as the extraction method, the type of solvent used and sample concentration (Tyasrini, 2006). In addition, *Candida albicans* fungi are a type of fungi that have virulent pathogenic causes and have morphological and tissue adhesion abilities so that they are more resistant and resistant to the administration of antifungal compounds contained in Agung Semeru banana peel extract of Lumajang variety.

Hand sanitizer is an antiseptic product that has an active ingredient in the form of 70% alcohol, so it can reduce the number of bacteria when used. As an antiseptic alcohol has the advantage of being volatile, so it does not require a long time to dry when applied on the hands. But this also becomes a weakness, because its effectiveness is only in the short term, so that bacteria can only be reduced in a short time after the use of antiseptics.

Hand sanitizer uses natural ingredients of Agung Semeru banana peel extract which has antifungal ability, as evidenced by the inhibition zone diameter at the highest concentration treatment. The antifungal effect on the hand sanitizer is due to the banana peel containing phytochemical compounds, namely phenolic, terpenoids, saponins and alkaloids (Sari and David, 2017). The mechanism of action of saponins by reducing the surface tension of cell membranes resulting in increased cell permeability or leakage and causing intracellular compounds to come out (Saroja *et al.*, 2012).

In addition, Saponin is a secondary metabolic compound that has an antiseptic function so that it is capable of being an antibacterial. Saponin compounds will form complex compounds with cell membranes through hydrogen bonds, so that the permeability properties of cell walls can be destroyed and cause cell death (Nur, 2013). Research conducted by Dinastuti (2015) that banana peel extract contains several active compounds that can be useful as antifungal, where the active compounds can affect the structure and function of *Candida albicans* cells such as cell walls and cell membranes.

The mechanism of action of alkaloid compounds is by disturbing the peptidoglycan component in the cell so that the cell wall layer is not fully formed, causing cell death (Saraswati, 2015). Phenol compounds are a group of tannin compounds and have natural antimicrobial activities that work by interacting with microbial cells through absorption processes that involve hydrogen bonds so that they can disrupt the mechanism of action of active transport on cells (Saefudin *et al.*, 2011).

The inhibitory response based on the inhibitory category according to Greenwood (1995 in Fitri

2010) is as follows: Inhibition zone diameter ≤ 10 mm is said to not inhibit the growth of test microbes (N), diameter 11-15 mm categorized as weak (W), diameter 16-20 mm as moderate (M) and diameter > 20 mm as strong (S). Based on the calculation results that Agung Semeru banana peel extract has a strong ability (S) in inhibiting the growth of *Candida albicans* (table 1), Based on the above explanation it can be concluded that the effectiveness test of Agung Semeru banana peel extract of Lumajang variety as a natural hand sanitizer is proven to be really true to inhibit the *Candida albicans*.

CONCLUSION

Utilization of Agung Semeru Banana peel Extract as a natural basic material hand sanitizer showed that no significant difference between treatments in inhibiting the growth of fungi *Candida albicans*, but a concentration of 7% ($1,315 \pm 0.0035^c$) showed better results compared to a concentration of 1% (1.2 ± 0.0^a); concentration of 5% ($1,208 \pm 0,0023^b$) and 0% ($1.2 \pm 0,0^a$).

ACKNOWLEDGEMENT

A big thank you to the researchers and IKIP PGRI Jember, especially FP. MIPA is given the opportunity to carry out research at the Laboratory of Biology FP. MIPA IKIP PGRI Jember, as well as all parties who helped in completing this research.

REFERENCE

¹⁶ Dinastutie, R. (2015). Uji efektifitas antifungal ekstrak kulit pisang kepok (*Musa acuminata x balbisiana*) mentah terhadap pertumbuhan

Candida albicans secara in vitro (Doctoral dissertation, Universitas Brawijaya).

¹² Dyer, D. L., Shinder, A., & Shinder, F. (2000). Alcohol-free instant hand sanitizer reduces elementary school illness absenteeism. *FAMILY MEDICINE-KANSAS CITY-*, 32(9), 633-638.

³ Ehiowemwenguan, G., Emoghenc, A.O Inctianbor, J.E. 2014. "Antibacterial and Phytochemical Analisis of Banana Fruit Peel, *IQSR Journal of Pharmacy*. Vol 4. Page: 18-25.

²¹ Fadhilah, N. L., & Rahayu, T. (2017). Potensi Pelepah Daun Pisang Kepok Sebagai Hand Sanitizer Alami (Doctoral dissertation, Universitas Muhammadiyah Surakarta).

¹⁴ Febriani, T. H. (2014). Uji Daya Antifungi Jus Buah Pare (*Momordica charantia* L) Terhadap Daya Hambat Pertumbuhan *Candida Albicans* Secara in vitro (Doctoral dissertation, Universitas Muhammadiyah Surakarta).

¹¹ Fitri, L. (2010). The effect of *Michelia alba* Bark Extract to The Growth of *Salmonella typhi* and *Candida albicans*. *Jurnal Natural*, 10(1), 27-30.

¹⁰ Hamzah, M. (2007). *Dermatoterapi Ilmu Penyakit Kulit dan Kelamin*, Edisi ke-5. FKUI, Jakarta, 342-52.

² Hasyim, N., Pare, K. L., Junaid, I., & Kurniati, A. (2012). Formulasi dan Uji Efektivitas Gel Luka Bakar Ekstrak Daun Cocor Bebek (*Kalanchoe pinnata* L.) Pada Kelinci (*Oryctolagus cuniculus*). *Majalah Farmasi dan Farmakologi*, 16(2), 89-94.

Hawkins, B. (2011). *Text book of gynaecology*. Edisi ke-15.

¹ Kholifah, S. N., Sari, D. N. R., & Anitasari, S. D. (2018). Pengaruh Tingkat Kematangan Dan

- Konsentrasi Ekstrak Kulit Pisang Agung Semeru Terhadap *Staphylococcus aureus*. *Bio-ma: Jurnal Biologi dan Pembelajaran Biologi*, 3(1).
- 20 Kusumaningtyas, E. (2015). Mekanisme Infeksi *Candida albicans* pada permukaan sel. Prosiding Lokakarya Nasional Penyakit Zoonosis.
- Manan, A., & Kharisma, A. (2012). Kelimpahan Bakteri *Vibrio* sp. pada Air Pembesaran Udang *Vannamei* (*Litopenaeus vannamei*) sebagai Deteksi Dini Serangan Penyakit Vibriosis [The Abundance Of *Vibrio* sp. Bacteria On Enlargement Water Of *Litopenaeus vannamei* As The Early Detection Of Vibriosis]. *Jurnal Ilmiah Perikanan dan Kelautan*, 4(2), 128-134.
- 8 Nur, J et al. (2013). "Bioaktivitas getah pelepah pisang ambon *Musa paradisiacal* var *Sapientum* terhadap Pertumbuhan Bakteri *Staphylococcus aureus*, *Pseudomonas aeruginosa* dan *Escherichia coli*". Fakultas Biologi, Universitas Hasanuddin.
- Prahardini, P. E. R., & Telp, M. (2018). Karakterisasi varietas unggul pisang mas kirana dan agung semeru di kabupaten lumajang. *Buletin Plasma Nutfah Vol 16 No.2* (online)
- 10 Rachmayanti, R. D. (2013). Penggunaan media panggung boneka dalam pendidikan personal hygiene cuci tangan menggunakan sabun di air mengalir. *Jurnal Promosi Kesehatan*, 1(1), 1-9.
- 4 Radji, M., Suryadi, H., & Ariyanti, D. (2012). Uji efektivitas antimikroba beberapa merek dagang pembersih tangan antiseptik. *Pharmaceutical Sciences and Research (PSR)*, 4(1), 1-6.
- 2 Retnosari, D. (2007). Pengaruh Substitusi Tepung Ikan oleh Tepung Belatung Terhadap Pertumbuhan Benih Ikan Nila. *Oreochromis niloticus*). Skripsi.
- 6 Rosenthal, V. D., Guzman, S., & Safdar, N. (2005). Reduction in nosocomial infection with improved hand hygiene in intensive care units of a tertiary care hospital in Argentina. *American journal of infection control*, 33(7), 392-397.
- Rostinawati, T. (2009). Aktivitas Antibakteri Ekstrak Etanol Bunga Rosella (*Hibiscus Sabdariffa* L.) Terhadap *Escherichia Coli*, *Salmonella Typhi* Dan *Staphylococcus Aureus* Dengan Metode Difusi Agar. Fakultas Farmasi. Bandung: Universitas Padjadjaran.
- 1 Sari, D. N. R. (2017). Analisis fitokimia ekstrak kulit pisang agung semeru dan mas kirana. *Bio-ma: Jurnal Biologi dan Pembelajaran Biologi*, 2(2).
- Sari, D. N. R., Susilo, D. K., Zakiyah, S. Z., Khoiriyah, W., & Kholifah, S. N. (2018). Formulasi Krim Ekstrak Kulit Pisang Agung Semeru dan Pisang Mas Kirana Varietas Lumajang. *Jurnal Biota*, 4(1), 24-28.
- 13 Saroja, M., Santhi, R., & Annapoorani, S. (2012). Wound healing activity of flavonoid fraction of *Cynodon dactylon* in Swiss albino mice. *International Research Journal of Pharmacy*, 3(2), 230-231.
- 7 Saraswati, F. N. (2015). Uji Aktivitas Antibakteri Ekstrak Etanol 96% Limbah Kulit Pisang Kepok Kuning (*Musa balbisiana*) Terhadap Bakteri Penyebab Jerawat (*Staphylococcus epidermidis*, *Staphylococcus aureus*, dan *Propionibacterium acne*). UIN Syarif Hi-dayatullah Jakarta.

Saifudin, A. Rahayu dan Teruna. (2011). Standar-
isasi Bahan Obat Alam. Yogyakarta : Graha
Media

15 Tyasrini, E., Winata, T. dan Susantina. (2006). Hub-
ungan antara sifat dan metabolit *Candida* sp.
dengan patogenesis kandidiasis. JKM. 6(1):
52-67.

1 Zakiya, S. Z., Sari, D. N. R., & Al Habib, I. M.
(2018, February). UJI DAYA HAMBAT
KOMBINASI EKSTRAK KULIT PISANG
AGUNG SEMERU (*Musa paradisiaca* L.)
DAN PISANG MAS KIRANA (*Musa acu-
minata* L.) TERHADAP *Candida albicans*. In
Prosiding Seminar Nasional SIMBIO-SIS
(Vol. 2).

9 Zubier, F., Bramono, K., Widaty, S., Nilasari, H.,
Louisa, M., & Rosana, Y. (2010). Efikasi
sabun ekstrak sirih merah dalam mengurangi
gejala keputihan fisiologis. *Majalah Kedokter-
an Indonesia*, 60(1).

UTILIZATION OF AGUNG SEMERU BANANA PEEL EXTRACT AS NATURAL HAND SANITIZER

ORIGINALITY REPORT

17%

SIMILARITY INDEX

16%

INTERNET SOURCES

9%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1	jurnal.radenfatah.ac.id Internet Source	3%
2	media.neliti.com Internet Source	1%
3	eprints.ums.ac.id Internet Source	1%
4	eprints.umm.ac.id Internet Source	1%
5	Esti Harpeni, Supono Supono, Dwi Risca Septiani. "THE GROWTH KINETICS OF Bacillus sp. D2.2 AT DIFFERENT pH AND SALINITY", AQUASAINS, 2019 Publication	1%
6	Submitted to Western Governors University Student Paper	1%
7	digilib.uinsby.ac.id Internet Source	1%

journals.ums.ac.id

8	Internet Source	1%
9	id.123dok.com Internet Source	1%
10	jurnal.unimus.ac.id Internet Source	1%
11	jbioua.fmipa.unand.ac.id Internet Source	1%
12	krex.k-state.edu Internet Source	1%
13	es.scribd.com Internet Source	1%
14	fr.scribd.com Internet Source	1%
15	jurnal.ar-raniry.ac.id Internet Source	1%
16	repository.ub.ac.id Internet Source	1%
17	Rahmayani Notanubun, Ritha L Karuwal. "HUBUNGAN KEKERABATAN FENETIK VARIETAS PISANG (Musa sp.) DI PULAU AMBON", BIOPENDIX: Jurnal Biologi, Pendidikan dan Terapan, 2014 Publication	<1%

18 Andrew P. Golin, Dexter Choi, Aziz Ghahary. "Hand Sanitizers: A Review of Ingredients, Mechanisms of Action, Modes of Delivery, and Efficacy Against Coronaviruses", American Journal of Infection Control, 2020
Publication <1%

19 www.neliti.com
Internet Source <1%

20 Submitted to Sriwijaya University
Student Paper <1%

21 eprints.poltekkesjogja.ac.id
Internet Source <1%

22 Dwi Nur Rikhmasari. "PERBANDINGAN KEMAMPUAN EKSTRAK KULIT PISANG AGUNG SEMERU DAN PISANG MAS KIRANA VARIETAS LUMAJANG DALAM MENGHAMBAT PERTUMBUHAN Candida albicans", Florea : Jurnal Biologi dan Pembelajarannya, 2017
Publication <1%

23 fmipa.unsyiah.ac.id
Internet Source <1%

24 core.ac.uk
Internet Source <1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

UTILIZATION OF AGUNG SEMERU BANANA PEEL EXTRACT AS NATURAL HAND SANITIZER

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
